Original Research Article

Histopathological spectrum of upper gastrointestinal endoscopic biopsies of neoplastic lesions: A retrospective study

Tarun Mittal¹, Niranjan J^{2,*}, Prathima S¹

¹Dept. of Pathology, Vydehi Institute of Medical Sciences and Research Centre, Bengaluru, Karnataka, India

²Dept. of Pathology, East Point Medical College, Bidrahalli, Karnataka, India

ARTICLE INFO

Article history:

Received 20-04-2020

Accepted 19-11-2020

Available online 18-12-2020

Keywords:

Endoscopy

Upper GI tract

Neoplastic lesions

ABSTRACT

Introduction: Upper gastrointestinal tract disorders are one of the commonly seen entities in the clinical practice. Gastric carcinoma is the second most common cancer while oesophageal carcinoma is the sixth leading cause of the death worldwide. Histopathological study has been done to make a specific diagnosis or to follow the evolution of particular disease.

Aims and Objectives: To study histopathological distribution of various types of neoplastic upper GI lesions.

Materials and Methods: A retrospective study was conducted on 1,800 upper gastrointestinal biopsies performed at our hospital from May 2015 to April 2018. The biopsies were formalin fixed, and embedded in paraffin wax, sections were taken, stained with Haematoxylin & Eosin and examined

Results: The present study included 1800 patients, out of which 486 (27%) cases were found to be neoplastic. The mostly commonly encountered neoplastic lesions were -well differentiated Adenocarcinoma (8.5%), followed by poorly differentiated Adenocarcinoma (6%) and adeno-squamous carcinoma. The male to female ratio was 1.5:1.

Conclusion: Upper gastrointestinal endoscopy is safe, cost effective, relatively less invasive procedure and its co-ordination with histopathological examination is the gold standard for early detection of GI tract lesions especially neoplastic lesions and thus aiding in early treatment and management.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Gastrointestinal tract is one of the commonest site for various lesions with high degree of mortality and morbidity. Further endoscopic biopsy is commonly performed procedure for variety of benign and malignant lesions.¹ Gastric carcinoma is the second most common carcinoma and is the leading cause of death in the current scenario.²

As per Indian National Cancer Register, oesophageal and gastric cancer are the commonest cancers found in men.³ Histopathological identification of gastrointestinal biopsies at the earlier stages allows an early intervention and prompt treatment of the disease.⁴

Introduction of endoscopes in 1960 has greatly improved the diagnostic facility. Tissue specimen can be taken from the lesion under direct vision using endoscopic biopsy forceps.¹ Upper Gastrointestinal endoscopy is the most sensitive and specific method for the detection of various neoplasms specially esophageal, gastric and duodenal neoplasm.⁵

Endoscopic biopsies detects lesion at an early stages and helps in prevention in progress of various lesions to carcinoma.⁶

2. Aims and Objectives

1. To study histopathological spectrum of various type of neoplastic upper gastrointestinal lesions

* Corresponding author.

E-mail address: tarunmittal90@gmail.com (Niranjan J).

- To study the age distribution, gender distribution and overall frequency of the lesion

3. Materials and Methods

It is a retrospective study conducted in the department of pathology, VIMS&RC, Bangalore on 1,800 upper gastrointestinal biopsies, performed at Vydehi hospital during a 36 months period from May 2015 to April 2018, out of which 486 (27%) cases were found to be neoplastic and were reviewed retrospectively.

Biopsies slides stained with H&E and Giemsa were retrieved from archival material and were studied in detail with respect to nature of the lesion, age distribution, gender distribution, and histopathological features.

3.1. Inclusion Criteria

- All endoscopic biopsies from upper gastrointestinal tract.

3.2. Exclusion Criteria

- All lesions of mouth and pharynx.
- Resection specimens not included
- Post treatment
- All duodenal biopsies beyond the second part of the duodenum.

4. Observations and Results

The present study includes 486 (27%) cases of upper gastrointestinal neoplastic biopsies between the period May 2015 to April 2018.

The following results were observed.

Table 1: Distribution of Lesions

Nature of lesion	No. of cases	Percentage (%)
Non-neoplastic	1,314	73
Neoplastic	486	27
Total	1800	100

Table 2: Gender wise distribution of neoplastic lesions

Gender	Neoplastic
Male	279
Female	207
Total	486

Out of 1,800 upper gastrointestinal endoscopic biopsy samples, that were studied during the period of three years, 486 (27%) were neoplastic. There were 279 males and 207 females making the male to female ratio of 1.4:1. The highest incidence was seen between 4th to 6th decade, while lowest incidence was seen in patients after 8th decade of life. The commonest site for gastric biopsy was pylorus followed

Table 3: Age Distribution

Age (Years)	Frequency	Percentage (%)
0-20	18	3.70
21-40	138	28.39
41-60	236	48.55
61-80	91	18.72
80+	3	0.64
Total	486	100

Table 4: Frequency and Percentage of Neoplastic Lesions

Histopathological Diagnosis	Frequency	Percentage (%)
Adeno Squamous Carcinoma	29	5.96
GIST	17	3.49
Squamous Papilloma	6	1.23
Well Differentiated Squamous Cell Carcinoma	44	9.05
Moderately Differentiated Squamous cell carcinoma	76	15.63
Poorly Differentiated Squamous cell carcinoma	30	6.17
Well Differentiated Adenocarcinoma	112	23.04
Moderately Differentiated Adenocarcinoma	51	10.49
Poorly Differentiated Adenocarcinoma	80	16.46
Adenocarcinoma - Signet Ring Type	34	6.99
Non-Hodgkin's Lymphoma - Stomach	2	0.41
Leiomyoma-Esophagus	5	1.02

Fig. 1: Non-Hodgkin's Lymphoma Stomach H&E Stain (10X)

Fig. 2: Signet Ring Adenocarcinoma Stomach H&E Stain (10X)

Fig. 4: Squamous papilloma H&E Stain (10X)

Fig. 3: Poorly differentiated Adenocarcinoma H&E Stain (10X)

Fig. 5: Gastro-intestinal Stromal Tumour (GIST) H&E Stain (40X)

by body, cardia and fundus.

Out of 486 neoplastic lesion, well differentiated adenocarcinoma is most common followed by poorly differentiated adenocarcinoma and the least common finding was non-Hodgkin's lymphoma stomach.

5. Discussion

In the present study, most common site for gastro intestinal biopsy is stomach followed by esophagus and duodenum. Endoscopic biopsy provides important clues that helps in diagnosis of various lesions.⁶ Clinical and endoscopic information is the fundamental part of adequacy and interpretation of biopsy is strongly affected by these factors.⁶ Out of 486 patients, with neoplastic endoscopic biopsies, males were predominantly affected i.e. 58% while rest 42% were females. This was also proved by another study done by Shennak MM et al⁷ over a period of 10 years.

The male: female ratio was 1.80:1. The ratio is suggestive of male predominance, mostly because of increased exposure of males to more risk factors, same established by the study done by JC Paymaster et al.⁸ In the present study, the highest incidence of upper gastrointestinal endoscopic lesions are seen between 41-60 years which comprises of 48.55% of total number of biopsies, similar to study conducted by Qureshi et al⁹ with 72 endoscopic biopsy samples over a period of 2 years. The oldest patient was 89 years old and the youngest was 9 years old. The age related difference could be due to the variations in risk factors among the different age groups. The incidence of neoplastic lesions was common in 5th decade with male predominance. Among the neoplastic lesion, stomach was most common site followed by esophagus.

Smoking, alcohol, dietary factors and social habits have been proposed as risk factor for gastric cancer.¹⁰ The incidence of gastric cancer is higher as compared to esophageal cancer. The highest incidence of neoplastic lesion in the present study was seen between 4th to 6th decade. This observation was similar to study carried out by Qureshi et al⁹ on 13,881 endoscopic biopsies over a period of 10 years.

Squamous dysplasia is a precursor lesion to squamous cell carcinoma. About 30% of dysplastic cases progress to carcinoma.¹¹ Obesity, alcohol consumption, smoking and reflux are the risk factors for adenocarcinoma esophagus, and is having bad prognosis and prognosis become worst in cases of signet ring and mucinous variant of adenocarcinoma.¹²

In our study, the most commonly encountered neoplastic lesion was well differentiated Adeno carcinoma (20.37%) followed by poorly differentiated Adeno carcinoma (14.61%) and Adeno-squamous carcinoma (14.2%).

6. Conclusion

In our study, commonest site for endoscopic biopsies was stomach, with a total number of 486(27%) neoplastic cases. The most commonly affected age group is between 40-60 years. Well differentiated Adeno carcinoma was a common malignancy followed by poorly differentiated Adenocarcinoma.

Upper gastrointestinal endoscopy is safe, simple, cost-effective, well tolerated and relatively less invasive procedure which permits exact diagnosis and helps in prompt treatment. Although histopathology is the gold standard, but biopsy provides a great opportunity to establish a definite/conclusive diagnosis and thus aiding in early treatment and management.

There were no cases of duodenal malignancy in our study.

7. Source of Funding

No financial support was received for the work within this manuscript.

8. Conflict of Interest

The authors declare they have no conflict of interest.

References

1. Krishnappa R, Horakerappa MS, Ali K, Gouri M. A study on histopathological spectrum of upper gastrointestinal tract endoscopic biopsies. *Int J Med Res Health Sci*. 2013;2(3):418–24. doi:10.5958/j.2319-5886.2.3.073.
2. Enzinger PC, Mayer RJ. Esophageal cancer. *N Engl J Med*. 2003;349:2241–52.
3. *National Cancer Registry Programme First All India Report*. 2001;1.
4. Hussain SI, Ruby R, Gulshan A, Ambreen B. Clinico histopathological study of upper Gastrointestinal tract endoscopic biopsies. *Int J Cur Res Rev*. 2015;7(16):78–85.
5. Aparajita A, Mohanty RC, Sahu AA. Histomorphological study of upper GI endoscopic biopsies. *Int J Health Sci Res*. 2016;6(12):59–64.
6. Suvakovic Z, Bramble MG, Jones R, Wilson C, Idle N, Ryott J, et al. Improving the detection rate of early gastric cancer requires more than open access gastroscopy: a five year study. *Gut*. 1997;41(3):308–13. doi:10.1136/gut.41.3.308.
7. Shennak MM, Tarawneh MS, Al-Sheikh TM. Upper Gastrointestinal Diseases in Symptomatic Jordanians: A Prospective Endoscopic Study. *Ann Saudi Med*. 1997;17(4):471–4. doi:10.5144/0256-4947.1997.471.
8. Paymaster JC, Sanghvi LD, Ganghadaran P. Cancer of gastrointestinal tract in western India. *Cancer*. 1968;21:279–87.
9. Qureshi NA, Hallissey MT, Fielding JW. Outcome of index upper gastrointestinal endoscopy in patients presenting with dysphagia in a tertiary care hospital-A 10 years review. *BMC Gastroenterol*. 2007;7(1):43. doi:10.1186/1471-230x-7-43.
10. Gajalakshmi V, Swaminathan R, Shanta V. An independent survey to assess completeness of Registration: Population based cancer registry. *Asian Pac J Cancer Prev*. 2001;2:179–83.
11. Boffetta P, Hashibe M. Alcohol and cancer. *Lancet Oncol*. 2006;7(2):149–56. doi:10.1016/s1470-2045(06)70577-0.
12. Howlader N, Noone AM, Krapcho M, Miller D, Bishop K, Kosary CL, et al. SEER data submission. In: SEER Cancer Statistics Review, 1975-2014, National Cancer Institute; 1976. Available from: https://seer.cancer.gov/csr/1975_2014/.

Author biography

Tarun Mittal, Post Graduate

Niranjan J, Professor

Prathima S, Professor and HOD

Cite this article: Mittal T, Niranjan J, Prathima S. Histopathological spectrum of upper gastrointestinal endoscopic biopsies of neoplastic lesions: A retrospective study. *IP J Diagn Pathol Oncol* 2020;5(4):410-414.