Original Research Article

To study and analyze hematological parameters in anaemia in males

Dupinder Kaur^{1,*}, Pooja Agarwal¹¹Dept. of Pathology, Shri Ram Murti Smarak Institute Of Medical Sciences, Bareilly, Uttar Pradesh, India

ARTICLE INFO

Article history:

Received 09-01-2021

Accepted 10-03-2021

Available online 14-06-2021

Keywords:

hematological

CBC

smear

males & anaemia

ABSTRACT

Background & Method: Hematological parameters like Hb (haemoglobin), TC (total count), DC (differential count), PCV (packed cell volume), MCV (mean red cell corpuscular volume) done in the automated cell counter and peripheral smear findings were studied.

Study Designed: Observational study.

Results: Out of 250 cases, 192 i.e. 76.8% cases showed microcytic hypochromic anaemia, 30 cases i.e. 12% had normocytic hypochromic anaemia, 27 cases i.e. 10.6% had normocytic normochromic anaemia and dimorphic anaemia was seen in 02 cases i.e. 0.6% cases. Out of 250 cases, 193 i.e. 77.2% cases showed microcytosis maximally in 0-5 years age group and 57 cases i.e. 22.8% had normocytic picture. The distribution of peripheral smear (RBC size) finding with age varied significantly (p value <0.05).

Conclusion: Iron deficiency is almost universal when dealing with this magnitude of anaemia. However, clinically speaking, many technical experts believe that to differentiate severe anaemia, a screening for other causes is desirable, all males are recommended to be screened. In the present study of pediatric cases 0-5 years age group males were most affected and prevalence was more in males as compared to females and the predominant morphological pattern was microcytic hypochromic anaemia.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Hemolytic weakness is characterized as abbreviated red platelet endurance because of either inherent deformities in the red platelets or outside factors.¹ By far most of erythrocyte problems that happen in the pediatric age bunch result from anomalies inside the red platelets (intra corpuscular defects).² Intra-corpuscular imperfections are quite often hereditarily decided, while, the irregularities inside the plasma are generally obtained. Periodically in G6PD inadequacy, an acquired irregularity of erythrocytes prompts hemolysis just when exceptional conditions exist in the plasma like the presence of specific medications, synthetics or other molecules.³

Among hemolytic anaemias Thalassemia major, Hereditary spherocytosis, G6 PD lack and procured causes like diseases and falciparum intestinal sickness can cause

extreme anaemia.⁴

The early history of megaloblastic weakness is the historical backdrop of malignant iron deficiency. The principal case was accounted for in 1822. In 1849, Addison depicted a patient with deadly idiopathic sickliness. For a long time, malignant paleness was secretly alluded to as "Addisonian vindictive weakness".⁵ Ehrlich presented the expression "megaloblast" to portray the enormous nucleated erythroid antecedents found in vindictive anaemia.⁶

Megaloblastic morphology might be found in various conditions. In guys, it results from a lack of folic corrosive or nutrient B12 or both. Vit. B12 and folic corrosive are the cofactors which are fundamental for the union of nucleoproteins and consequently, their lack brings about damaged union of DNA and RNA which prompts incapable erythropoiesis and diminished life expectancy of RBCs.⁷

* Corresponding author.

E-mail address: dupindercaur@gmail.com (D. Kaur).

2. Materials and Methods

Our study is a hospital based study done in a tertiary care health center. Study duration 2014 Study was conducted at 2016 Hematological parameters like Hb (haemoglobin), TC (total count), DC (differential count), PCV (packed cell volume), MCV (mean red cell corpuscular volume) done in the automated cell counter and peripheral smear findings were studied.

2.1. Inclusion criteria

1. Below 15 years male
2. WHO criteria for anaemia
3. Males with hemoglobin < 11gm/dl

2.2. Exclusion criteria

1. Males above 15 year of age
2. Hemoglobin above 11 gm/dl
3. Hb Electrophoresis, wherever applicable

3. Results

Table 1: Distribution of the various morphological types of anaemia

Morphological Types	Number	Percentage
Microcytic Hypochromic Anaemia	192	76.8
Normocytic Hypochromic Anaemia	30	12
Normocytic Normochromic Anaemia	27	10.6
Dimorphic Anemia	2	0.6

Table 1 Out of 250 cases, 192 i.e. 76.8% cases showed microcytic hypochromic anaemia, 30 cases i.e. 12% had normocytic hypochromic anaemia, 27 cases i.e. 10.6% had normocytic normochromic anaemia and dimorphic anaemia was seen in 02 cases i.e. 0.6% cases.

Table 2 The distribution of peripheral smear (RBC chromasia) with age did not differ significantly as p value >0.05.

Table 3 Out of 250 cases, 193 i.e. 77.2% cases showed microcytosis maximally in 0-5 years age group and 57 cases i.e. 22.8% had normocytic picture. The distribution of peripheral smear (RBC size) finding with age varied significantly (p value <0.05).

4. Discussion

In the current investigation, all the cases having MCV <59fl showed 100% microcytic RBCs, none was normocytic on fringe smear assessment. MCV between 59-80fl showed microcytic cells in 79% cases and normocytic cells in 21% cases.⁸ MCV >80 fl showed 96% cases with normocytic

Table 2: Age wise distribution of cases according to peripheral smear (RBC Chromasia) finding

Peripheral Smear (RBC Chromasia) Finding	Age Group		11-15 years		Total	P Value
	No.	%	No.	%		
Dimorphic	02	1.55	01	1.80	02	p-value=
Hypochromic	114	87.69	46	83.63	222	0.396, Pearson
Normochromic	14	10.76	08	14.57	26	Chi-Square =
Total	130	100	55	100	250	3.146

Table 3: Age wise distribution of cases according to peripheral smear (RBC size) finding

Peripheral Smear (RBC Size) Finding	0-5 years		Age Group 6-10 years		11-15 years		Total		P Value
	No.	%	No.	%	No.	%	No.	%	
Microcytic	104	80.62	51	78.46	37	67.72	193	77.20	P-Value = 0.047, Pearson Chi-Square = 8.166
Normocytic	26	19.37	15	21.53	18	32.72	57	22.80	
Total	130	100	66	100	55	100	250	100	

cells and 4% cases with microcytic cells.

MCH <20 pg showed 100% cases having hypochromasia. MCH between 20-27 pg showed 90% cases having hypochromic cells alongside 9.47% cases having normochromic cells and 0.38% cases had dimorphic populace which was independently included as far as hypochromasia.⁹ MCH>27pg showed 84.85% cases having normochromic cells followed by 9.09% cases having hypochromic cells and 6.06% cases had dimorphic population.¹⁰ The p-esteem for this relationship between's MCH finding with chromasia was 0.000 for example p-esteem <0.005.

From both the examinations it was noticed that RBC records are identified with size and chromasia of RBC.¹¹ This importance is credited to acceptable quality control of the cell counter in our institution and normal adjustment of the same.¹² Be that as it may, the MCV is an incredibly helpful incentive in arrangement of anaemias¹³ however the MCH regularly don't add critical and clinically important data. Since MCH assume a significant part in research center quality control on the grounds that the qualities will stay stable for a given example over time.¹⁴

5. Conclusion

Iron deficiency is almost universal when dealing with this magnitude of anaemia. However, clinically speaking, many technical experts believe that to differentiate severe anaemia, a screening for other causes is desirable, all males are recommended to be screened. In the present study of pediatric cases 0-5 year's age group males were most affected and prevalence was more in males as compared to females and the predominant morphological pattern was microcytic hypochromic anaemia.

6. Source of Funding

No financial support was received for the work within this manuscript.

7. Conflicts of Interest

There are no conflicts of interest.

References

1. Sharada B. Iron deficiency and iron deficiency anaemia. *Recent Adv Malass.* 1995;5:205-33.
2. Saloojee H, Pettifor JM. Iron deficiency and impaired child development. *BMJ.* 2001;323(7326):1377-8. doi:10.1136/bmj.323.7326.1377.
3. Israels MCG. The pathological significance of the megaloblast. *J Pathol.* 1939;49(1):231-40.
4. Watkins D, Whitehead VM, Rosenblatt DS. Megaloblastic Anaemia. In: Nathan DG, Horkin S, Ginsburg D, Look TA, Fisher D, editors. 7th Edn.. In: Nathan and Oski's Hematology of infancy and childhood. Philadelphia: Elsevier; 2008. p. 468-78.
5. Bourner G, Dhaliwal J, Sumner J. Performance Evaluation of the Latest Fully Automated Hematology Analyzers in a Large, Commercial Laboratory Setting: A 4-Way, Side-by-Side Study. *Lab Hematol.* 2005;11(4):285-97. doi:10.1532/lh96.05036.
6. Tefferi A, Hanson CA, Inwards DJ. How to Interpret and Pursue an Abnormal Complete Blood Cell Count in Adults. *Mayo Clin Proc.* 2005;80(7):923-36. doi:10.4065/80.7.923.
7. CLSI. validation, verification and quality assurance of automated hematology analyzers. In: 2nd Edn. Approved Standard. CLSI document H26-A2. Wayne, PA: Clinical and Laboratory Standards Institute; 2010.
8. Firkin F, Chesterman C, Penington D, Rush B. de Gruchy's clinical haematology in medical practice. In: 5th Edn.; 2011. p. 24-5.
9. Rehema H, Simbauranga E, Kamugish A, Hokororo BR, Kidenya J. Makani Prevalence and factors associated with severe anaemia amongst under-five males hospitalized at. *Bugando Med Centre.* 2015;15:13.
10. Pandey A, Pandey M, Dombale VD, Sawke GK, Sawke N. Haematological Profile of Anaemia in Males of Age 1-12 Years in Southern Rural India; People's. *J Scientific Res.* 2016;9(1):48-52.
11. Muthuraman M, Sintha M. A cross sectional analysis of Anaemia in males population in a tertiary care hospital Madurai. *Med Pulse Int J Pathol.* 2017;1(2):42-7.
12. Rathna S, Venkatraman J, Govindaraj AS, Patil P. Study of Morphological Pattern of Anaemia in Males. *J Evol Med Dent Sci.* 2014;3(27):7540-3.
13. Gao W, Yan H, Wang D, Dang S, Pei L. Severity of Anaemia among males under 36 months old in rural Western China. *J Pone.* 2013;8(4):1-6.
14. Poornima S, Balaji PAR, Varne SRR, Jayashree K, Saba F. Anemia among hospitalized children at a multispecialty hospital, Bangalore (Karnataka), India. *J Family Med Prim Care.* 2014;3(1):48-53. doi:10.4103/2249-4863.130275.

Author biography

Dupinder Kaur, Resident

Pooja Agarwal, Associate Professor

Cite this article: Kaur D, Agarwal P. To study and analyze hematological parameters in anaemia in males. *IP J Diagn Pathol Oncol* 2021;6(2):132-134.