Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Journal of Diagnostic Pathology and Oncology

Journal homepage: <https://www.jdpo.org/>

Original Research Article

Role of needle biopsy in early diagnosis of lung lesions in tertiary care hospital

S.H Khaparde¹, Sneha B Patrike^{1,*}, S.D Deshmukh¹, B.B Shinde¹¹Dept. of Pathology, DVVPFS Medical College under MUHS, Nashik, Maharashtra, India

ARTICLE INFO

Article history:

Received 12-07-2021

Accepted 02-09-2021

Available online 13-09-2021

Keywords:

Adenocarcinoma

Computed tomography

Lung biopsy

Malignant

ABSTRACT

Background: Lung cancer is one of the most common cancer worldwide and the leading cause of cancer-related death. Needle biopsies of lung have been used for diagnosis of lung lesions. They are done CT guided or thorascopic guided procedures in our institution.

Aims: The aim is to study role of needle biopsy in early diagnosis of lung lesions and incidence of lung lesions in needle biopsies in our hospital with respect to age and sex.

Materials and Methods: A retrospective type of study. The histopathological records of our Hospital were reviewed for needle biopsies of lung lesions between January 2019 to December 2019 (One year). 35 cases noted during this period.

Results: There were a total of 35 cases during the study period of one year. The specimens received were linear bits and average size was 0.5cm in length. Non-neoplastic lesions were eight (22.85 %) and neoplasms contributed to 15 cases (42.85%) along with inadequate biopsies (20 %) and normal lung biopsies (14.28 %). Maximum number of cases were reported in the age group of 51-70 years (28 cases) with male preponderance. In all age groups, malignant lesions were more common in which, Adenocarcinoma was predominant.

Conclusion: To conclude, in our study malignant neoplasms predominated over non neoplastic lesions. The most common histological type encountered in this study was an Adenocarcinoma.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](#), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

The Lungs are involved in variety of inflammatory, neoplastic and various other lesions.¹ Around the world millions of people suffer from various preventable chronic respiratory diseases.² A range of conditions that involve the lung parenchyma may be associated with inflammation, fibrosis or granulomatous reactions.³ Lung cancer is a major public health problem. Prompt pathological investigation and diagnosis is essential as most of the clinic-radiological findings in various respiratory illnesses are nonspecific and are essential to improve patient survival, to avoid the rapid progression of the disease and to spare the patient from

more invasive procedures^{4,5} Therefore, it is most important to determine the leading causes of death due to pulmonary diseases to establish correct prophylactic actions & is the least expensive strategy for prevention of further pulmonary irreversible dysfunction.⁶

It is the most commonly diagnosed cancer worldwide accounting for 1.61 million (12.7%) of the total cancer patients.⁷ Every year more than 1.6 million deaths occur due to Lung cancer, more than deaths occurring due to breast, colon and prostate cancers combined.⁸ All over world it accounts for 13% of all new cancer cases and 19% of cancer related deaths.⁹ Lung is also one of the most common metastatic sites of many primary cancers.¹⁰ According to studies, lung carcinoma is a frequent finding amongst all the chest diseases.¹¹

* Corresponding author.

E-mail address: s.patrike1989@gmail.com (S. B. Patrike).

In males it is the 2nd most common cancer, while in females 6th most common cancer & the mortality rate is 4th after breast, cervix, and oral cavity-lip cancer.¹²

Non-specific symptoms of lung cancer and high prevalence of tuberculosis in our country which often leads to misdiagnosis and delayed diagnosis of lung cancer and also many patients present with various co-morbidities which adversely affect diagnosis and prognosis of the disease.¹³

Cigarette smoking is strongly related to Small cell lung carcinoma which rapidly metastasizes to many sites within the body and are most often discovered very late after they have spread extensively throughout body.¹⁴ Other various factors affecting include occupational exposures, air pollution and old pulmonary scars.¹⁵

Smoking tobacco (both cigarettes & beedis) is the principal risk factor for lung cancer in Indian men; however, among Indian women, the association with smoking is not that much strong, suggesting that there must be other risk factors other than smoking in causation of lung cancer.¹⁶ Despite numerous advances in recent years in terms of diagnostic methods, molecular changes, and therapeutic interventions, the outcomes of the lung cancer patients remain poor; hence, a better understanding of the risk factors may impact the preventive measures to be implemented at the community level.

Tissue biopsy is often required to differentiate primary lung lesion from metastasis, benign lesions from malignant lesions and further classify a primary lung cancer for treatment.¹⁵

Small lung biopsies are the most common and the first lung sample obtained when a radiological abnormality is detected, tissue diagnosis is required.¹⁷ Small lung biopsies are the primary method of diagnosis for the majority of lung cancer patients because 70% of lung cancers are unresectable as patients present in very advanced stages.¹⁸ Needle biopsies have been documented to be most effective for diagnosis of peripheral lung lesions including both non-neoplastic and neoplastic lesions.¹⁰

After introduction of lung cancer screening tests more patient majority in early stages of the disease are diagnosed based on small specimens. Furthermore, these specimens are needed for an accurate histopathological classification as well as for molecular testing.^{19–21}

2. Aims and Objectives

Available data will be analyzed,

1. To study role of needle biopsy in early diagnosis of lung lesions.
2. To study incidence of lung lesions in needle biopsies in our hospital with respect to age and sex.
3. Also to classify tumors according to WHO guidelines whenever possible to render specific diagnosis.

3. Materials and methods

The data for this study has been taken from medical record department (MRD) between January 2019 to December 2019 (1 year). There were 35 cases during this period.

3.1. Inclusion criteria

All the guided biopsies of lung were included in this study irrespective of them being non-neoplastic and neoplastic, benign or malignant nature including primary and metastases.

3.2. Exclusion criteria

Lobectomy specimen and pneumonectomy specimen, Pleural & mediastinal biopsies were excluded from this study.

The specimen received were linear bits and average size was 0.5cm in length. The specimen was fixed in 10% Neutral buffered formalin and subjected to routine tissue processing then paraffin embedded sections were made. This was followed by H&E staining and preparation of slide. Information including age, gender and final clinical diagnosis were collected from the medical record sections. Detailed clinical history and radiological findings were also obtained.

4. Results

There were total of 35 cases during the study period of one year. Non-neoplastic lesions were eight (22.85 %) and neoplasms contributed to 15 cases (42.85%) along with Inadequate biopsies seven cases (20 %) and normal lung biopsies were 5 (14.28 %). [Table 1]

Maximum number of cases were reported in the age group of 51-70 years (28 cases) followed by five cases in 31-50 years age group and only two cases in 0 – 29 years age group with male preponderance. In age groups of 51-70 years and 31-50 years, malignant lesions were more common in which Adenocarcinoma was predominant. [Table 4]

Out of eight non neoplastic lesions encountered, most common being inflammatory (37.5 %) followed by Pneumonia (25%) and others (25 %), fibrosis (12.5 %). The Others category includes one case of Hydatid cyst and one case of congenital cystic adenomatoid malformation. [Table 1]

All 15 neoplastic lesions encountered in our study were malignant. Among them histologically, Adenocarcinoma was most common followed by squamous cell carcinoma. Adenocarcinoma accounted in seven cases i.e 46.66 %. In two cases of Adenocarcinoma, histologically Lepidic pattern was seen. Squamous cell carcinoma cases were four (26.66 %).

Table 1: Non neoplastic lesions of needle lung biopsies

Diagnosis	Number of cases
Inflammatory	3 (37.5 %)
Pneumonia	2 (25 %)
Others	2 (25 %)
Fibrosis	1 (12.5 %)
Total	8 cases

Table 2: Malignant lesions of needle lung biopsies

Diagnosis	Number of cases
Adenocarcinoma	7 (46.66 %)
Squamous cell carcinoma	4 (26.66 %)
Small cell carcinoma	1 (6.66 %)
No subtype (Positive for malignancy)	2 (13.33 %)
MALT lymphoma	1 (6.66 %)
Total	15 cases

Table 3: Comparison of Malignancies with various studies

Histologic types	Present study	Agarwal A et al ²²	Krishnamurthy A et al ⁷	Sundaram V et al ²³
Adenocarcinoma	46.66 %	29.3 %	42.6 %	43.33 %
Squamous cell carcinoma	26.66 %	21.88 %	15.8 %	31.67 %
Small cell carcinoma	6.66 %	12.8 %	13.1 %	10 %
Undifferentiated carcinoma	-	24 %	6.9 %	11.67 %
Metastasis	-	6.4 %	-	-

Table 4: Demographic profile and distribution of histological subtypes of lung carcinoma in various studies

Study	Mean age	Male to female ratio	Squamous cell carcinoma	Adenocarcinoma	Small cell carcinoma	Other
Present study	58	3.9 : 1	26.66 %	46.66 %	6.66 %	20 %
Krishnamurthy et al ⁷	56	3.5 : 1	15.8 %	42.6 %	13.1 %	28.3 %
Malik et al ²⁴	55	4.6 : 1	25.1 %	41 %	14.8 %	19.1 %
Noronha et al ²⁵	56	3.5 : 1	24.1 %	40.3 %	8 %	27.6 %

There were two cases (13.33 %) in which subtyping was not possible due to small biopsies but they were malignant i.e. Positive for malignancy. One case of MALT Lymphoma was encountered accounting for 6.66 % of all malignant lesions. [Table 2]

Low power 10x

High power 40x

Fig. 1: ADENOCARCINOMA (Moderately differentiated)

Low power 10x

High power 40x

Fig. 2: Lepidic cell carcinoma

Fig. 3: Papillary carcinoma

Fig. 4: Spindle cell variant

5. Discussion

During the study period of One year there were found a total of 35 cases, maximum number of cases were seen in 5th to 7th decade, whereas in studies done by Agarwal A et al²² and Malik PS et al²⁴ the common age group affected was 5th to 6th decade.

The male to female ratio in the present study was 5.2 : 1 and in studies done by Malik PS et al,²⁴ Agarwal et al²² and Noronha et al²⁶ the ratios were 4.6:1 , 4.7:1 and 3.5 :1 respectively. Malignancies were more common than non-neoplastic lesions accounting to 15 cases out of 35 in our study which was in contrast to a study done by Kulshrestha R et al²⁶ where non neoplastic lesions were more common accounting for 506 cases out of 916 cases.

The malignant lung lesions were seen commonly in age group of 51-70 years and mean age was 58 years which was comparable to various studies by Noronha V et al²⁵ Dey A et al,²⁷ Koul PA et al²⁸ where 56, 57.9, 57.6 respectively were the mean age.

The male to female ratio of malignancies was 3.9:1 with male preponderance which was very similar to studies done by Kumar et al,²⁹ Pandhi N et al,³⁰ Dhandapani et al³¹ , Gupta et al.³² This showed that in India still females have lower prevalence of smoking.³³

The most common malignancy seen in our study was Adenocarcinoma (seven cases, 46.66%) which was much more similar to studies done by Krishnamurthy et al,⁷ Agarwal A et al,²² Sundaram et al²⁴ and Malik PS et al²⁵ while in some studies by Kulshrestha R et al,²⁶ Gupta R et al,³² Sarfraz S et al³³ and Mandal SK et al³⁴ the most common malignancy encountered was Squamous cell carcinoma.

Similar to studies done by Krishnamurthy et al,⁷ Agarwal A et al,²² Malik PS et al,²⁴ the second common malignancy seen was Squamous cell carcinoma (Four cases).

Among the non-neoplastic lesions of lung, in our study majority cases were inflammatory lesions (three cases, 37.5%). This was followed by two cases (25%) of Pneumonia.

Our study finding were in contrast to the study done by Kulshrestha R et al²⁶ in which most of the cases were granulomatous lesions of lung (30.2%) which was followed by inflammatory lesions and fibrosis.

According to most of studies in India, squamous cell carcinoma is the most dominant subtype, but many of the recent studies from India show changing trends of carcinoma, reporting Adenocarcinoma as the predominant subtype.³⁵

6. Conclusion

To conclude, malignant lung lesions were predominant than non -neoplastic lesions of lung in our study. In our study, the most common histological type encountered was an Adenocarcinoma. In the present study, inflammatory lesions was the most common non- neoplastic condition followed by pneumonia.

7. Conflict of Interest

There is no potential conflict of interests related to the exclusive nature of this paper.

8. Source of Funding

No financial support was received for the work on this manuscript.

References

1. Bal MS, Sethi PS, Suri AK, Bodal VK, Kaur G. Histopathological pattern in lung autopsies. *JPAFMAT*. 2008;8(2):29–31.
2. Hall JE. Guyton and Hall Textbook of Medical Physiology. In: and others, editor. 13th Edn. Elsevier: Saunders; 2015.
3. Kumar V, Abbas A, Aster J. Robbins & Cotran Pathologic Basis of Disease. In: 9th Edn.. vol. 2. Elsevier; 2014.
4. Chauhan G, Agrawal M, Thakkar N, Parghi B. Spectrum of histopathological lesions in lung autopsy. *J Res Med Den Sci*. 2015;3(2):109–12.
5. Jameson JL, Fauci A, Kasper D, Hauser S, Longo D, Loscalzo J, et al. Harrison's principles of internal medicine. In: 19th Edn.. vol. 2. India: McGraw Hill; 2015.

6. Ghosal R, Kloer P, Lewis KE. A review of novel biological tools used in screening for the early detection of lung cancer. *Postgraduate Med J*. 2009;85(1005):358-63.
7. Behera D. Lung cancer in India. *Indian J Chest Dis Allied Sci*. 2004;46(4):269-81.
8. Krishnamurthy A, Vijayalakshmi R, Gadigi V, Ranganathan R, Sagar T. The relevance of "Nonsmoking- associated lung cancer" in India. A single-center experience. *Indian J Cancer*. 2012;49(1):82-8.
9. Siegel R, Naishadham D, Jemal A. Cancer statistics. *Cancer J Clin Sci*. 2015;62(1):10-29. doi:10.3322/caac.20138.
10. Liang L, Jing L, Buryanek J, Zhang S. CT-guided core needle biopsy of peripheral lung lesions with onsite adequate valuation: Review of 215 cases. *J Cytol Histol*. 2014;4:1-5. doi:10.4172/2157-7099.S4-018.
11. Dubey N, Arti J, Varudkar HG, Agrawat JC, Deepali B, Sunil M, et al. A clinico-pathological profile of primary lung cancer patients presenting in a rural medical college of Central India. *Panacea J Med Sci*. 2015;5(3):124-9.
12. International Agency for Research on Cancer. Cancer Fact Sheets. [Last accessed on 2018 Nov 29]; 2018. Available from: <http://www.gco.iarc.fr/today/data/factsheets/populations/356‑>.
13. Behera D. Epidemiology of lung cancer a Global and Indian perspective. *JACM*. 2012;13(2):131-7.
14. Inamura K. Lung cancer: Understanding its molecular pathology and the 2015 WHO classification. *Front Oncol*. 2017;7:1-6. doi:10.3389/fonc.2017.00193.
15. Garima B, Renuka IV, Ramya P, Prasad D. Pravallika Uppala and Prathyusha. Histopathological Study of Guided Lung Biopsies. *Ann Pathol Lab Med*. 2020;7(4):180-5.
16. Noronha V, Pinninti R, Patil VM, Joshi A, Prabhash K. Lung cancer in the Indian subcontinent. *South Asian J Cancer*. 2016;5(3):95-103.
17. Mukhopadhyay S, Mehta AC. Utility of Core needle biopsies and transbronchial biopsies for diagnosing Nonneoplastic lung diseases. *Arch Pathol Lab Med*. 2018;142(9):1054-68.
18. Travis WD, Brambilla E, Müller-Hermelink HK, Harris CC. Pathology and Genetics: Tumors of the Lung, Pleura, Thymus and Heart. France: IARC; Lyon; 2004.
19. Travis WD, Brambilla E, Noguchi M. The new IASLC/ATS/ERS international multidisciplinary lung adenocarcinoma classification. *J Thoracic Oncol*. 2011;6(5):244-85. doi:10.3978/j.issn.2072-1439.2014.01.27.
20. Lindeman NI, Cagle PT, Beasley MB. Molecular testing guideline for selection of lung cancer patients for EGFR and ALK tyrosine kinase inhibitors: guideline from the College of American Pathologists, International Association for the Study of Lung Cancer, and Association for Molecular Pathology. *J Thorac Oncol*. 2013;8(7):823-59. doi:10.1097/JTO.0b013e318290868f.
21. Leighl NB, Rekhman N, Biermann WA. Molecular testing for selection of patients with lung cancer for epidermal growth factor receptor and anaplastic lymphoma kinase tyrosine kinase inhibitors: American Society of Clinical Oncology endorsement of the College of American Pathologists/International Association for the study of lung cancer/association for molecular pathology guideline. *J Clin Oncol*. 2014;32(32):3673-9. doi:10.1200/JCO.2014.57.3055.
22. Agarwal A, Tandon R, Singh L, Kumar P, Pant H. Clinical profile of lung cancer in a tertiary care teaching hospital in North India with special reference to acceptance and outcome of treatment. *J Pulmon*. 2018;2:4-8.
23. Sundaram V, Sanyal N. Clinicopathological profile of bronchogenic carcinoma in a tertiary care hospital in eastern part of India. *Clin Cancer Investig J*. 2014;3(3):220-4.
24. Malik PS, Sharma MC, Mohanti BK, Shukla NK, Deo S, Mohan A, et al. Clinico-pathological profile of lung cancer at AIIMS: A changing paradigm in India. *Asian Pac J Cancer Prev*. 2013;14(1):489-94. doi:10.7314/apjcp.2013.14.1.489.
25. Noronha V, Dikshit R, Raut N, Joshi A, Pramesh CS, George K, et al. Epidemiology of lung cancer in India: Focus on the differences between non-smokers and smokers: A single-centre experience. *Indian J Cancer*. 2012;49(1):74-81. doi:10.4103/0019-509X.98925.
26. Kulshrestha R, Menon BK, Vijayan VK. Role of a pattern based approach in interpretation of transbronchoscopic lung biopsy and its clinical implications. *Indian J Chest Dis Allied Sci*. 2012;54(1):9-17.
27. Dey A, Biswas D, Saha SK, Kundu S, Sengupta A. Comparison study of clinicoradiological profile of primary lung cancer cases: An Eastern India experience. *Indian J Cancer*. 2012;49(1):89-95. doi:10.4103/0019-509X.98930.
28. Koul PA, Kaul SK, Sheikh MM, Tasleem RA, Shah A. Lung cancer in the Kashmir valley. *Lung India*. 2010;27(3):131-7. doi:10.4103/0970-2113.68309.
29. Kumar M, Sharma DK, Garg M, Jain P. Clinicopathological profile of lung cancer - changing trends in India. *Int J Res Med*. 2016;5(2):57-62.
30. Pandhi N, Malhotra B, Kajal N, Prabhudesai RR, Nagaraja CL, Mahajan N, et al. Clinicopathological profile of patients with lung cancer visiting Chest and TB Hospital Amritsar. *Sch J App Med Sci*. 2015;3(2):802-9.
31. Dhandapani S, Srinivasan A, Rajgopalan R, Chellamuthu S, Rajkumar A, Palaniswamy P, et al. Clinicopathological profile of lung cancer patients in teaching hospital in South India. *J Cardiothorac Med*. 2016;4:442-3.
32. Gupta R, Chowdary I, Singh P. Clinical, radiological and histological profile of primary lung carcinomas. *JK Sci*. 2015;17(3):146-51.
33. Sarfraz S, Gupta R, Bhardwaj S. Histopathological Patterns of endobronchial lung biopsy specimen in lung cancer along with clinico-radiological correlation. *Int J Contemp Med Res*. 2018;5(11):1-5.
34. Mandal SK, Singh TT, Sharma TD, Amrithalingam V. Clinico-pathology of lung cancer in a regional cancer centre in Northeastern India. *Asian Pac J Cancer Prev*. 2013;14(12):7277-81. doi:10.7314/apjcp.2013.14.12.7277.
35. Prasad R, James P, Kesarwani V, Gupta R, Pant MC, Chaturvedi A, et al. Clinico-pathological study of bronchogenic carcinoma. *Respirology*. 2004;9:557-60.

Author biography

S.H Khaparde, Professor and HOD

Sneha B Patrike, Junior Resident

S.D Deshmukh, Professor

B.B Shinde, Professor

Cite this article: Khaparde SH, Patrike SB, Deshmukh SD, Shinde BB. Role of needle biopsy in early diagnosis of lung lesions in tertiary care hospital. *IP J Diagn Pathol Oncol* 2021;6(3):170-174.