Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Journal of Diagnostic Pathology and Oncology

Journal homepage: <https://www.jdpo.org/>

Case Report

Rhinoscleroma – A case report with review of literature

Meenakshi Mohapatro¹, Lipika Behera¹, Shushruta Mohanty^{1,*},
Swetambari Acharya¹, Swayamprava Pradhan¹

¹Dept. of Pathology, M.K.C.G. Medical College, Berhampur, Odisha, India

ARTICLE INFO

Article history:

Received 25-08-2021

Accepted 01-09-2021

Available online 13-09-2021

Keywords:

Rhinoscleroma

mikulicz cells

chronic granulomatous infection

ABSTRACT

Rhinoscleroma is a chronic granulomatous disease caused by gram negative rod shaped bacteria *Klebsiella rhinoscleromatis*. Most common sites of predilection include the nasal cavity and nasopharynx, but there are reports of its occurrence at other sites like the larynx, trachea, bronchi, middle ear, and orbit. We are reporting here a case of rhinoscleroma in a middle aged female patient involving the nasal cavity along with a brief review of literature.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](#), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Rhinoscleroma is rare chronic infectious granulomatous process involving the upper respiratory tract caused by *Klebsiella rhinoscleromatis*¹ which is a short encapsulated gram negative rod shaped bacilli. It is endemic in Middle east, central and south America and is sporadic in other parts of the world.²⁻⁶ It has a female preponderance affecting older age groups in 3rd to 6th decade. Main route of transmission is by direct inhalation of nasal droplets or indirect contact with nasal secretions of infected person.⁷ Although it has affinity for nasal mucosa it is known to occur at other sites like nasopharynx (18-43%), larynx(15-40%), trachea (12%) and bronchi (2-7%).⁸ 10 % of rhinoscleroma cases are also known to occur in the lip.⁹ Here we are reporting a case of rhinoscleroma that was clinically suspected as inverted papilloma.

2. Case Report

A 34 year female presented to ENT out patient department with a reddish swelling inside right nostrils since 3

months. It was insidious in onset and gradually progressive. She complained of long standing history of sneezing, nasal itching with nasal obstruction and on and off headache with running nose for which she consulted a local physician who prescribed her with medications. Her symptoms reduced a bit however the mass didn't subside on taking those medications so she was then referred to our institution with a suspicion of the mass being malignant. On examination the swelling was non tender, soft to firm in consistency and did not bleed on touch. She was planned for incisional biopsy for which she was advised to do a panel of routine blood tests, reports of which were apparently normal. Punch biopsies were taken and was send to our department for histopathological analysis. Grossly we received bits of irregular grayish black tissue altogether measuring 3x2 cm. Slides were stained with routine H&E. All slides were examined under light microscopy. H&E stains showed stratified squamous epithelium with underlying subepithelium showing groups of large vacuolated histiocytes with single or multiple cytoplasmic vacuoles and small centrally to peripherally located nuclei (mikulicz cells), fine nuclear chromatin and indistinct nucleoli admixed with chronic inflammatory cells

* Corresponding author.

E-mail address: sushruta.mohanty@gmail.com (S. Mohanty).

lymphocytes and plasma cells along with Russel bodies (Figures 1, 2, 3 and 4). Giemsa stained demonstrated the intracellular bacilli within the clear cytoplasmic vacuoles (Figure 5). Thus Final diagnosis of rhinoscleroma was clinched based on histopathology and special stains.

Fig. 1: Scanner view 40x showing epithelial lining with subepithelium showing vacuolated cells.

Fig. 2: LP 100x view showing the foamy cells in the subepithelium having single to multiple cytoplasmic vacuoles and small centrally to peripherally located nuclei (mikulicz cells), fine nuclear chromatin and indistinct nucleoli.

3. Discussion

Rhinoscleroma was first described by Hebra in the year 1870. The disease predominantly occurs in middle aged woman, maintaining a poor hygienic and nutritional status. Humans are the only host for *K. rhinoscleromatis*.

Fig. 3: LP100x showing mikulicz cells admixed with lymphoplasmacytic infiltrate

Fig. 4: LP100x Showing lymphoplasmacytic infiltrate in the subepithelium with mikulicz cells.

The exact pathogenesis of the disease as to how mikulicz cell is formed and eventually rhinoscleroma formation is not clear. *Klebsiella rhinoscleromatis* like *Klebsiella pneumoniae* has a mucopolysaccharide capsule that contributes to inhibition of phagocytosis thus facilitating its intracellular survival.¹⁰ It is also postulated that altered immune response with impaired cellular immunity is thought to play a role in the pathogenesis of the disease and an alteration in CD4:CD8 ratio is responsible for chronicity of the disease.¹¹ Rhinoscleroma usually occurs at the epithelial transition zones, especially at the junction where stratified squamous epithelium of the vestibule meets ciliary epithelium of the nose.¹²

Fig. 5: Oil immersion 1000xGiemsa stain –Showing presence of intracellular gram negative bacilli within the vacuolated cells (Mikulicz cells).

Fig. 6: LP 100x- Fite Faraco staining showing absence of lepra bacilli within the vacuolated cells.

Clinically the patient may present with any of the three stages of disease progression. Three stages described in rhinoscleroma are: catarrhal (rhinitic), granulomatous (proliferative or nodular), and sclerotic (cicatricial or fibrotic).¹ Patients with initial catarrhal phase present with nonspecific rhinitis that evolves into chronic foul smelling purulent discharge along with dry crusting of mucous membrane, and nasal obstruction. On histopathology, it shows a nonspecific subepithelial infiltrate of polymorphonuclear leucocytes with granulation tissue.² The clinical features are more prominent in

granulomatous phase. In this phase, patient develops granulomatous nodular rubbery mass(lump) which leads to nasal obstruction, loss of smell and may lead to nasal enlargement with nasal deformity. The dense infiltration of lymphocytes, plasma cells, Russell bodies, and pathognomonic large Mikulicz cells are the diagnostic histopathological changes seen in granulomatous stage.² If left untreated patient goes into sclerotic phase. Permanent complications are the features of sclerotic phase which includes nasal deformities, dysphonia, dysphasia, stridor and blocked airways.¹ In sclerotic phase, extensive fibrosis and less inflammatory cell infiltration are the main histopathological findings.^{1,2}

Histopathological differential diagnosis for rhinoscleroma is quite broad and includes both neoplastic and nonneoplastic disease states. Given the morphologic appearance of Mikulicz cells, one must consider histiocytic lesion, such as Rosai-Dorfman disease as a differential diagnosis. Rosai-Dorfman disease classically presents with massive, painless cervical lymphadenopathy with extranodal disease most commonly affecting the skin and central nervous system. Histology reveals histiocytes and emperipolesis with immunohistochemistry positivity for S100 and CD68 but negative results for CD1a.¹³ In younger patients, disseminated juvenile xanthogranuloma and Langerhans cell histiocytosis may be considered. Juvenile xanthogranuloma presents typically with a dermal lesion, showing histiocytes, mixed inflammation, and Touton giant cells and can be disseminated in rare cases. Immunohistochemistry is positive for CD68 and negative for S100 and CD1a.¹⁴ Langerhans cell histiocytosis commonly involves the head and neck region, lacks Touton giant cells, has a prominent eosinophilic infiltrate, and is strongly positive for CD1a.¹⁵ Under Infectious category most important differential includes leprosy that resembles histologically and is characterised by presence of diffuse arrangement of foamy macrophages (virchow cells) that contains plenty of lepra bacilli within them. They are usually highlighted by doing special stains like (fite faracco stain) that was negative in our case [Figure 6] and ruled out the possibility of leprosy.

4. Conclusion

The diagnosis of rhinoscleroma thus relies on the adequate correlation of clinical, histopathological, and bacterial identification by the use of special stains. Around 50% to 60% of patients shows positive cultures for *K* rhinoscleromatis on blood or MacConkey agar who presents in the granulomatous stage.¹⁶ Identification of *Klebsiella* species can also be accomplished with the help of 16s rRNA gene sequencing.¹⁷ Main stay of treatment of Rhinoscleroma includes a combination of antibiotics, surgical debridement and repair; however, the patient needs to be on routine follow up as the recurrence rates are very

high. Our patient was started on ciprofloxacin 500 mg and doxycycline 100 mg twice daily for 4 months and showed good response to treatment with no clinical relapse even after 6 months of follow up.

5. Conflict of Interest

There is no potential conflict of interests related to the exclusive nature of this paper.

6. Source of Funding

No financial support was received for the work on this manuscript.

References

1. Mukara BK, Munyarugamba P, Dazert S, Löhler J. Rhinoscleroma: A case series report and review of the literature. *Eur Arch Otorhinolaryngol.* 2014;271(7):1851–6. doi:10.1007/s00405-013-2649-z.
2. Moraes MA, Magalhães AV, Marinho LC, Azevedo AE, Carneiro FP, It R, et al. Rhinoscleroma causing severe bilateral nasal obstruction. *Braz J Infect Dis.* 2010;14(2):190–2. doi:10.1590/s1413-86702010000200015.
3. Hart CA, Rao SK. Rhinoscleroma. *J Med Microbiol.* 2000;49(5):395–6. doi:10.1099/0022-1317-49-5-395.
4. Muzyka MM, Gubina KM. Problems of the epidemiology of scleroma. I. Geographical distribution of scleroma. *J Hyg Epidemiol Microbiol Immunol.* 1971;15(3):233–42.
5. Muzyka MM, Gubina KM. Problems of the epidemiology of scleroma. 2. Some aspects of the problem of endemic focus formation. *J Hyg Epidemiol Microbiol Immunol.* 1972;16(1):8–20.
6. Chan TV, Spiegel JH. Klebsiella rhinoscleromatis of the membranous nasal septum. *J Laryngol Otol.* 2007;121(10):998–1002. doi:10.1017/S0022215107006421.
7. Inamadar AC, Palit A, Kulkarni NH, Guggarigoudar SP, Yelikar BR. Nodulo ulcerative lesions over the nose. *Indian J Dermatol Venereol Leprol.* 2004;70(3):197–8.
8. Bailhache A, Dehesdin D, François A, Marie JP, Choussy O. Rhinoscleroma of the sinuses. *Rhinology.* 2008;46(4):338–41.
9. Miller RH, Shulman JB, Canalis RF, Ward PH. Klebsiella rhinoscleromatis: A clinical and pathogenic enigma. *Otolaryngol Head Neck Surg.* 1979;87(2):212–21. doi:10.1177/019459987908700211.
10. Hoffmann EO, Loose LD, Harkin JC. The Mikulicz cell in rhinoscleroma: light, fluorescent and electron microscopic studies. *Am J Pathol.* 1973;73(1):47–58.
11. Posthumus JS, Schneyer MS. Pathology quiz case 1. Rhinoscleroma. *Arch Otolaryngol Head Neck Surg.* 2011;137(10):1043–5. doi:10.1001/archoto.2011.144-a.
12. Razek A, A A. Imaging of scleroma in the head and neck. *Br J Radiol.* 2012;85(1020):1551–5. doi:10.1259/bjr/15189057.
13. Dalia S, Sagatys E, Sokol L, Kubal T. Rosai-Dorfman disease: tumor biology, clinical features, pathology, and treatment. *Cancer Control.* 2014;21(4):322–7.
14. Sanguenza OP, Salmon JK, White CR, Beckstead JH. Juvenile xanthogranuloma: a clinical, histopathologic and immunohistochemical study. *J Cutan Pathol.* 1995;22(4):327–35.
15. Buchmann L, Emami A, Wei JL. Primary head and neck Langerhans cell histiocytosis in children. *Otolaryngol Head Neck Surg.* 2006;135(2):312–7.
16. Ahmed AR, Badawy ZH, Mohamed IR, Abdelhameed WA. Rhinoscleroma: A detailed histopathological diagnostic insight. *Int J Clin Exp Pathol.* 2015;8(7):8438–45.
17. Brisse S, Grimont F, Grimont PAD. The genus Klebsiella. *Prokaryotes.* 2006;6:159–96.

Author biography

Meenakshi Mohapatro, Assistant Professor

Lipika Behera, Assistant Professor

Shushruta Mohanty, Assistant Professor

Swetambari Acharya, Post Graduate

Swayamprava Pradhan, Professor and HOD

Cite this article: Mohapatro M, Behera L, Mohanty S, Acharya S, Pradhan S. Rhinoscleroma – A case report with review of literature. *IP J Diagn Pathol Oncol* 2021;6(3):245-248.